



**Fungicide Resistance Management Strategies**  
**Developed by the CropLife Australia Fungicide Resistance Management Review Group**  
Valid as at 25 June 2015

**Crop(s): Barley**

**Pest(s): Powdery Mildew, Scald, Net blotch**

Resistance Management Strategy for:

**Group 11 + 3** (Quinone outside Inhibitor + DMI)

Treat as a **Group 11** fungicide for resistance management purposes.

1. Apply only as a protectant treatment – where no more than 5% leaf area infection evident anywhere in the canopy.
2. **DO NOT** apply more than two applications per growing season.
3. **DO NOT** apply consecutive sprays within a spray program.
4. **DO NOT** rotate with alternative DMI fungicides which are known to have compromised resistance status.
5. In high risk disease environments integrated management approaches should be used to reduce fungicide resistance risk including the removal of stubble, control of green bridge volunteers and the use of tolerant and resistant varieties.

Resistance Management Strategy for:

**Group 3** (DMI)

1. Apply only as a protectant treatment – where no more than 5% leaf area infection evident anywhere in the canopy.
2. Rotate with products which incorporate alternative modes of action partners with activity on powdery mildew where possible
3. **DO NOT** apply more than three applications per growing season. The total of three applications includes DMI's applied as **Group 11 + 3** co-formulations and in-furrow or seed treatments that have activity on foliar diseases. Combinations of in-furrow and seed treatment are counted as one application.
4. **DO NOT** rotate with alternative DMI fungicides which are known to have compromised resistance status.
5. In high risk disease environments integrated management approaches should be used to reduce fungicide resistance risk including the removal of stubble, control of green bridge volunteers and the use of tolerant and resistant varieties. This is particularly important for managing barley powdery mildew which has shifted in tolerance to DMI fungicides in some regions.

Resistance Management Strategy for:

**Group 7** (SDHI)

1. Apply only as a protectant treatment – where no more than 5% leaf area infection evident anywhere in the canopy.
2. Rotate with products which incorporate alternative modes of action partners with activity on powdery mildew where possible
3. **DO NOT** apply more than two applications per growing season. This includes in-furrow or seed treatments that have activity on foliar diseases. Combinations of in-furrow and seed treatment are counted as one application.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.

Phone: 02 6230 6399 Email: [info@croplife.org.au](mailto:info@croplife.org.au) Fax: 02 6230 6355 Website: [www.croplife.org.au](http://www.croplife.org.au)