



Fungicide Resistance Management Strategies
Developed by the CropLife Australia Fungicide Resistance Management Review Group
 Valid as at 25 June 2015

Crop(s): **Onion**

Pest(s): **Downy Mildew**

Resistance Management Strategy for:

Group 4 (Phenylamide); and
Group 40 (Carboxylic acid amide) fungicides

1. Start disease control early and maintain a regular program using a fungicide from groups other than **Group 4** or **40**.

2. When conditions favour disease development, **DO NOT** wait for disease to appear, but apply two consecutive sprays of a **Group 4** or **40** product at the interval recommended on the label. Then resume the program of sprays using products from a different group to the **Group 4** or **40** products just applied.

3. **DO NOT** apply more than four sprays of a **Group 4** or **40** product per season.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.
 Phone: 02 6230 6399 Email: info@croplife.org.au Fax: 02 6230 6355 Website: www.croplife.org.au