



---

**Fungicide Resistance Management Strategies**  
**Developed by the CropLife Australia Fungicide Resistance Management Review Group**  
Valid as at 25 June 2015

---

**Crop(s):**        **Ornamentals**

**Pest(s):**       **Grey Mould**

Resistance Management Strategy for:

**Group 2**            (Dicarboximide) fungicides; and  
**Group 9**            (Anilinopyrimidine)

1.     If three or fewer Botrytis fungicide sprays are applied per crop, use only one spray containing a **Group 9** fungicide. If four to six sprays are applied per crop, use a maximum of two sprays containing **Group 9** fungicides. If seven or more sprays are applied per crop use a maximum of three sprays containing **Group 9** fungicides.

Avoid applying more than two successive sprays of a **Group 9** fungicide, including from the end of one season to the next.

2.     **DO NOT** apply more than two consecutive sprays of a **Group 2** fungicide.

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.  
Phone: 02 6230 6399   Email: [info@croplife.org.au](mailto:info@croplife.org.au)   Fax: 02 6230 6355   Website: [www.croplife.org.au](http://www.croplife.org.au)