

Fungicide Resistance Management Strategies
Developed by the CropLife Australia Fungicide Resistance Management Review Group
Valid as at 25 June 2015

Crop(s): Pulse Crops

Disease (s): Grey Mould (Botrytis)

Resistance Management Strategy for

Group 1 (Methyl Benzimidazole Carbamates);
Group 2 (dicarboximide); and
Group M3 (dithiocarbamate) fungicides

1. **DO NOT** apply more than two consecutive **Group 1** sprays, including a seed treatment containing a **Group 1** fungicide. Apply no more than two consecutive sprays of a **Group 1** fungicide alone.
2. **DO NOT** apply more than two **Group 1** treatments (including seed treatment) in a season.
3. **DO NOT** apply more than two **Group 2** sprays in one season. Apply no more than two consecutive sprays of a **Group 2** fungicide.
4. **DO NOT** apply more than two **Group M3** treatments (including seed treatment) in a season. Apply no more than two consecutive sprays of a **Group M3** fungicide alone.
5. It is advisable to rotate groups between seasons.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.
Phone: 02 6230 6399 Email: info@croplife.org.au Fax: 02 6230 6355 Website: www.croplife.org.au