

---

**Insecticide Resistance Management Strategies**  
**Developed by the CropLife Australia Insecticide Resistance Management Review Group**  
Valid as at 25 June 2015.

---

**Crop(s) : Strawberries/ornamentals**

**Mite : Two Spotted Mite**

**Guidelines:**

1. Monitor mite activity and treat infestations before thresholds are reached, ie. spray earlier rather than later. Seek advice on local threshold levels.
2. Do not apply sequential applications of products from any one chemical group.
3. Preferably products with the same Mode of Action should not be used more than twice in a growing season
4. Incorporate the use of predatory mites for the control of this pest wherever possible.

**Insect : Western Flower Thrips**

For information refer Industry and Investment NSW website

<http://www.dpi.nsw.gov.au/agriculture/horticulture/pests-diseases-hort/multiple/thrips/wft-resistance>

---

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.

Phone: 02 6230 6399 Email: [info@croplife.org.au](mailto:info@croplife.org.au) Fax: 02 6230 6355 Website: [www.croplife.org.au](http://www.croplife.org.au)