



Fungicide Resistance Management Strategies
Developed by the CropLife Australia Fungicide Resistance Management Review Group
Valid as at 25 June 2015

Crop(s): Strawberry

Pest(s): Grey Mould (Botrytis)

Resistance Management Strategy for

Group 2	(Dicarboximide);
Group 7	SDHI (Succinate dehydrogenase inhibitors)
Group 9	(Anilinopyrimidine);
Group 12	(Phenylpyrroles); and
Group 17	(Hydroxylanilide) fungicides

1. Apply a program of protectant fungicides during flowering. If conditions favour disease development during this period, use a **Group 2, 9, 12 or 17** fungicide.
2. **DO NOT** apply more than two successive sprays of **Group 2, 9, 12 or 17** fungicides.
3. If applying **Group 7** fungicides, **DO NOT** apply more than two consecutive sprays before changing to another group. **DO NOT** apply more than three **Group 7** sprays per season. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another **Group 7**, including sprays in consecutive seasons.
4. If three or fewer Botrytis fungicide sprays are applied per crop, use only one spray containing a **Group 9** fungicide. If four to six sprays are applied per crop use a maximum of two sprays containing **Group 9** fungicides. If seven or more sprays are applied per crop use a maximum of three sprays containing **Group 9** fungicides.
5. If three or fewer Botrytis fungicide sprays are applied per crop, use only one spray containing a **Group 12** fungicide. If four to six sprays are applied per crop use a maximum of two sprays containing **Group 12** fungicides. If seven or more sprays are applied per crop use a maximum of three sprays containing **Group 12** fungicides.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use.

Advice given in this strategy is **valid as at 25 June 2015**. All previous versions of this strategy are now invalid.
Phone: 02 6230 6399 Email: info@croplife.org.au Fax: 02 6230 6355 Website: www.croplife.org.au