



Fungicide Resistance Management Strategies

*Developed by the CropLife Australia Fungicide Resistance Management Review Group
and industry researchers*

Crop(s): Stone Fruit

Pest(s): Blossom Blight and Brown Rot

Resistance Management Strategy for:

- | | |
|----------------|---|
| Group 2 | (Dicarboximide); |
| Group 3 | (DMI); |
| Group 7 | SDHI (Succinate dehydrogenase inhibitors) and |
| Group 9 | (Anilinopyrimidine) fungicides |

1. If applying **Group 2** or **Group 3** fungicides, **DO NOT** apply more than two consecutive sprays of fungicides from the same group before changing to another group.
2. **DO NOT** apply more than three sprays of a **Group 9** fungicide per season. If two or three consecutive sprays are applied, they must be followed by at least the same number of sprays from an alternative chemical group, including from one season to the next.
3. If applying **Group 7** fungicides, **DO NOT** apply more than two consecutive sprays before changing to another group. **DO NOT** apply more than three **Group 7** sprays per season. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another **Group 7**, including sprays in consecutive seasons.
4. A post-harvest treatment should also be counted as an application.
5. The last blossom blight spray and the first pre-harvest brown rot spray should be regarded as consecutive applications.
6. The spray program should be considered and the strategy applied on a whole-orchard basis.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is **valid as at 7 June 2017**.

All previous versions of this strategy are now invalid. Phone: 02 62732733 Email: info@croplife.org.au Website: www.croplife.org.au