

Crop(s): Almonds

Pest(s): Blossom Blight and Brown Rot

Resistance Management Strategy for:

Group 2	(Dicarboximide) fungicides;
Group 3	Demethylation inhibitor (DMI) fungicides;
Group 7	SDHI (Succinate dehydrogenase inhibitors)
Group 11	QoI (Quinone outside inhibitor) fungicides
Group 11+3	QoI + DMI fungicides
Group 11+7	QoI + SDHI fungicides

1. **DO NOT** apply consecutive sprays of solo products containing **Group 7**. Consecutive sprays include mixtures containing **Group 7**.
2. **DO NOT** apply consecutive sprays of solo products containing **Group 11**. Consecutive sprays include mixtures containing **Group 11**.
3. **DO NOT** apply more than three **Group 3**, **Group 7** or **Group 11** sprays per season (including mixtures of **Group 11+3** and **Group 11+7**).
4. **DO NOT** apply more than three **Group 2** sprays in one season. Apply no more than two consecutive sprays before changing to another group.
5. If applying **Group 7** or **Group 11** fungicides in mixtures, **DO NOT** apply more than two consecutive sprays before changing to another group.
6. Consecutive application includes from the end of one season to the start of the following season.
7. Rotate with products from **Groups M2, M3 and M4**.
8. The spray program should be considered and the strategy applied on a whole-orchard basis.