

Crop(s): Fruit (post-harvest treatment)

Pest(s): Post-harvest diseases

Resistance Management Strategy for:

Group 2	(Dicarboximide); and other “systemic” fungicides; and
Group 3	(DMI);
Group 9	(Anilinopyrimidine) fungicides
Group 12	(Phenylpyrroles) fungicides

1. For the last pre-harvest spray, use a fungicide with a different activity group to the fungicide planned for use as a post-harvest treatment.
2. Where alternatives are available, rotate to use as many different activity groups as possible.
3. **DO NOT** dispose of unused dip solutions as a spray to crops or orchards.
4. **DO NOT** dispose of unused dip solutions within or near the crop or orchard area.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator’s (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is **valid as at 22 June 2018**. All previous versions of this strategy are now invalid.

Phone: 02 62732733 Email: info@croplife.org.au Website: www.croplife.org.au