

**Crop(s):** Lettuce

**Pest(s):** Botrytis & Sclerotinia

Resistance Management Strategy for:

|                 |  |
|-----------------|--|
| <b>Group 2</b>  | (dicarboximide);                           |
| <b>Group 7</b>  | SDHI (Succinate dehydrogenase inhibitors); |
| <b>Group 9</b>  | (anilinopyrimidine);                       |
| <b>Group 11</b> | (Quinone outside inhibitor); and           |
| <b>Group 12</b> | (phenylpyrroles) fungicides                |

1. Apply **Group 2** fungicide as a seedling drench soon after emergence.
2. Apply a protectant fungicide as a high volume foliar spray before planting out, then a **Group 2** fungicide immediately after planting.
3. Maintain a cover with protectant fungicide sprays at 7-10 day intervals.
4. If weather conditions favour Botrytis infection, tank mix the protectant with a **Group 2** fungicide.
5. **DO NOT** apply **Group 2** fungicides more than four times per season, irrespective of the target disease.
6. **DO NOT** apply more the two **Group 9** or **Group 12** fungicides per crop.
7. **DO NOT** apply more than three **Group 7** or **Group 11** fungicides per crop and no more than two consecutive sprays per crop.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 22 June 2018. All previous versions of this strategy are now invalid.

Phone: 02 62732733 Email: [info@croplife.org.au](mailto:info@croplife.org.au) Website: [www.croplife.org.au](http://www.croplife.org.au)