

Crop(s): Tomato

Pest(s): Grey Mould

Resistance Management Strategy for:

Group 2 (Dicarboximide) fungicides; and
Group 7 SDHI (Succinate dehydrogenase inhibitors)

1. Tank mix **Group 2** fungicides with a protectant such as chlorothalonil. Avoid applying two **Group 2** fungicides in succession, unless tank mixed with a protectant.
2. **DO NOT** apply more than four **Group 2** sprays in a season.
3. If applying **Group 7** fungicides, **DO NOT** apply more than two consecutive sprays before changing to another group. **DO NOT** apply more than three **Group 7** sprays per season. If consecutive sprays are used, then use the same number of sprays of an alternative group before using another **Group 7**, including sprays in consecutive seasons.

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been taken with the information supplied in this document no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian regulator's (APVMA) product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is **valid as at 22 June 2018**. All previous versions of this strategy are now invalid.