

SPECIFIC GUIDELINES FOR GROUP L HERBICIDES

GROUP	L	HERBICIDE
-------	---	-----------

Moderate resistance risk

Globally herbicide resistance to the Group L herbicide mode of action has been confirmed and documented in more than 30 weed species across 16 countries

Group L resistance exists in Australia in annual ryegrass, and in 2 species of barley grass across more than 100 populations, blackberry nightshade, crowsfoot grass, capeweed, pennsylvanian cudweed, squirrel-tailed fescue (silver grass) and small square weed. Most instances have occurred in long-term lucerne stands treated regularly with a Group L herbicide but Group L resistant barley grass has also occurred in no-till situations.

The following factors are common to most cases of Group L resistance:

- A Group L herbicide is the major or only herbicide used;
- A Group L herbicide has been used for 12 – 15 years or more; and
- There has been minimal or no soil disturbance following application.

The risk of resistance to Group L herbicides is higher in minimum/zero tillage broadacre cropping. Other high resistance risk situations include: irrigated clover pivots, orchards, vineyards or pure lucerne stands where frequent applications of a Group L herbicide are made each season, cultivation is not used and there is reliance on a Group L herbicide alone for weed control.

To assist in delaying the onset of resistance, consider alternating Group L herbicides with herbicides from other modes of action. For example, Group N (e.g. glufosinate) or Group Q (e.g. amitrole) or Group M (e.g. glyphosate).

Below are strategies that address these high resistance risk situations to reduce the risk of Group L resistance developing.

Minimum/Zero Tillage

1. Rotate Group L herbicides with other knockdown herbicides with a different mode of action, such as Group M (e.g. glyphosate). A full label rate for the weed size targeted should be used for resistance management.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 10 June 2020. All previous versions of this strategy are now invalid.

2. Consider utilising the double knock technique¹ where glyphosate is sprayed first followed within 1 - 7 days by a paraquat application. A full label rate for the weed size targeted should be used for the paraquat application for resistance management.
3. Consider occasional mechanical cultivation to aid weed control.

Lucerne

1. If using a Group L herbicide for winter cleaning, where possible include another mode of action e.g. Group C.
2. Use alternative modes of action to selectively control grass and broadleaf weeds.
3. Rotate Group L herbicides with other knockdown herbicides with a different mode of action (such as Group M e.g. glyphosate) prior to sowing lucerne and prior to sowing future crops in that paddock.

Horticulture

1. Rotate Group L herbicides with other knockdown herbicides with a different mode of action, such as Group N (e.g. glufosinate), Group Q (e.g. amitrole) or Group M (e.g. glyphosate).
2. Where possible, use residual herbicides (that are effective on the same weeds as the Group L herbicides) where applicable either alone or in mixture with Group L herbicides.
3. Where possible use alternative modes of action to selectively control grass and broadleaf weeds.
4. Consider using the double knock technique where glyphosate is sprayed followed within 1-7 days by a paraquat application. A full label rate for the weed size targeted should be used for the paraquat application for resistance management.

The above recommendations should be incorporated into an Integrated Weed Management (IWM) program. In all cases try to ensure surviving weeds from any treatment do not set and shed viable seed. Keep to integrated strategies mentioned in this brochure including cultural weed control techniques to reduce the weed seedbank. Make sure you mix and rotate herbicides from different mode of action groups. Always consult the product label prior to use.

Chemical family	Active constituent (first registered trade name)
-----------------	--

¹ The double knock technique is defined as using a full cut cultivation OR the full label rate of a paraquat-based product (Group L) following the glyphosate (Group M) knockdown application

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 10 June 2020. All previous versions of this strategy are now invalid.

GROUP L

Inhibitors of photosynthesis at photosystem I via electron diversion (PSI inhibitors)

Bipyridyls	diquat (Reglone®, Spray Seed®*), paraquat (Alliance®*, Gramoxone®, Spray Seed®*)
------------	--

* This product contains more than one active constituent

Notes:

1. List of chemical families, approved active constituents and, in parenthesis, the trade name of the first registered product or successor. Refer to the APVMA website (www.apvma.gov.au) to obtain a complete list of registered products from the PUBCRIS database.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 10 June 2020. All previous versions of this strategy are now invalid.