

Crop(s): **Wheat**

Disease(s): **Yellow Spot, Powdery Mildew**

Resistance Management Strategy for:

Group 3	Demethylation Inhibitors (DMI);
Group 7	Succinate dehydrogenase Inhibitor (SDHI);
Group 7+3	SDHI + DMI;
Group 11+3	Quinone outside Inhibitor (QoI) + DMI; and
Group 11+4	QoI + Phenylamide.
Group 11+7+3	QoI+SDHI+DMI

1. Fungicides should be used as protectant treatments – where no more than 5% leaf area infection evident anywhere in the canopy. In high risk disease environments, integrated management approaches should be used to reduce fungicide resistance risk including the removal of stubble, control of green bridge volunteers and the use of tolerant and resistant varieties.
2. **Do not** apply more than two applications per growing season of **Group 11** or **7** containing products. This includes in-furrow or seed treatments that have activity on foliar diseases. Combinations of in-furrow and seed treatment are counted as one application.
3. **Do not** apply consecutive applications of **Group 11** containing products. This includes in-furrow i.e. If a **Group 11+4** fungicide has been used in-furrow at planting, the first foliar fungicide spray must not contain a **Group 11** fungicide.
4. If a **Group 7** seed treatment has been used with foliar activity (as determined by label claims), the first foliar fungicide applied should not contain a **Group 7** fungicide.
5. **Do not** apply more than one application of a **Group 7** seed treatment with foliar activity in any two consecutive growing seasons.
6. If a **Group 7** fungicide is being applied as a foliar spray, it must be in a co-formulation or in mixture with a registered mixing partner with a different mode of action, with no known resistance.
7. **Do not** apply more than three applications containing **Group 3** fungicides per growing season. This total of three applications includes DMIs applied as **Group 11+3** or **Group 11+7+3** co-formulations and in-furrow or seed treatments that have activity on foliar diseases. Combinations of in furrow and seed treatment are counted as one application.
8. Minimise use of **Group 3** fungicides which are known to have compromised resistance status.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 10 June 2020. All previous versions of this strategy are now invalid.