

**Crop(s): Cucurbits**

**Disease(s): Powdery mildew**

Resistance Management Strategy for:

<b>Group 3</b>	Demethylation inhibitors (DMI);
<b>Group 7</b>	Succinate dehydrogenase inhibitors (SDHI);
<b>Group 8</b>	Hydroxy-(2-amino-) pyrimidines;
<b>Group 11</b>	Quinone outside inhibitors (QoI);
<b>Group 11+7</b>	QoI + SDHI;
<b>Group 13</b>	Aza-naphthalenes;
<b>Group U6</b>	Phenyl-acetamide; and
<b>Group 50</b>	Actin disruptors (aryl-phenyl-ketones).

1. Start disease control early. **Do not** wait for powdery mildew to appear before spraying but start as soon as practicable after crop emergence.
2. Use protectant sprays in early crop growth. Apply protectant sprays up to the fruit set stage of the crop if the disease normally occurs during this period. If this schedule is interrupted (e.g. by rain) use a tank mix of protectant plus systemic before recommencing the protectant program.
3. After fruit set, use systemic fungicides in one or more of the following ways:
4. Tank mix systemic fungicides with a protectant fungicide **and** use fungicides from at least two different systemic activity groups per crop.
5. Alternate systemic fungicides with a protectant fungicide **and** use fungicides from at least two different systemic activity groups per crop.
6. Alternate systemic fungicides from at least three different activity groups per crop.
7. Apply **Group 11** fungicides preventatively.
8. Use a maximum of one **Group 11** containing spray out of every three fungicide applications.
9. **Do not** use consecutive applications of **Group 11** or **Group U6** fungicides.
10. **Do not** apply more than two **Group 11** (including **11+7**) or **Group U6** products per crop.
11. **Do not** apply more than three **Group 7** or **Group 13** products per crop and no more than two consecutive applications per year.

**Please note:**

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 16 July 2021. All previous versions of this strategy are now invalid.