

Crops(s) **Peanut**

Disease(s) **Leaf spot, rust, net blotch**

Resistance management strategy for:

Group 3	Demethylation inhibitors (DMI);
Group 11	Quinone outside Inhibitors (QoI); and
Group 3+11	DMI + QoI

Guidelines:

1. **Do not** apply more than three consecutive **Group 3** sprays alone, before changing to a fungicide of a different activity group.
2. Apply a maximum of five **Group 3** sprays per season.
3. **Do not** apply **Group 11** products as more than 50% of the total number of sprays in any one season, up to a maximum of three sprays of **Group 11**.
4. **Do not** apply more than two consecutive applications of a **Group 11** spray.
5. If two consecutive applications of a **Group 11** spray are applied, they must be followed by at least the same number of sprays from an alternative chemical group.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 30 June 2022. All previous versions of this strategy are now invalid.