

Crops(s) **Fruit (post-harvest treatment)**

Disease(s) **Post-harvest diseases**

Resistance management strategy for:

Group 2	Dicarboximide and other "systemic" fungicides;
Group 3	Demethylation inhibitors (DMI);
Group 3+12	DMI + Phenylpyrroles;
Group 9	Anilinopyrimidine;
Group 11+12	Quinone outside Inhibitors (QoI) + Phenylpyrroles; and
Group 12	Phenylpyrroles.

Guidelines:

1. For the last pre-harvest spray, use a fungicide with a different activity group to the fungicide planned for use as a post-harvest treatment.
2. Where alternatives are available, rotate to use as many different activity groups as possible.
3. **Do not** dispose of unused dip solutions as a spray to crops or orchards.
4. **Do not** dispose of unused dip solutions within or near the crop or orchard area.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 14 July 2023. All previous versions of this strategy are now invalid.