

**Crops(s)**            **Apple, Pear**

**Disease(s)**        **Apple and Pear scab**

Resistance management strategy for:

<b>Group 3</b>	Demethylation inhibitors (DMI);
<b>Group 7</b>	Succinate dehydrogenase inhibitors (SDHI);
<b>Group 7+11</b>	SDHI + QoI;
<b>Group 9</b>	Anilinopyrimidines (AP);
<b>Group 11</b>	Quinone outside inhibitors (QoI); and
<b>Group U12</b>	Guanidines.

### Guidelines:

1. Not all listed products are registered for both diseases. Ensure the product is registered for the disease being targeted.
2. **Do not** apply more than four sprays per season of **Group 3** fungicides as solo products; if more sprays containing **Group 3** are required apply as a tank mixture with a non-cross resistant fungicide.
3. **Do not** apply more than four sprays per season of products containing **Group 9**. In locations where resistance has been reported use a **Group 9** only in mixture with a registered, alternative mode of action for which resistance is not known.
4. **Do not** apply more than three sprays per season of **Group 7** or **Group 11** fungicides.
5. If two consecutive applications of **Group 7, 9** or **Group 11** fungicides are used, then they must be followed by at least the same number of applications of fungicide(s) from a different mode of action group(s) before the same group is used again, either in the current or following season.
6. When mixtures are used for resistance management, applied as tank mix or as a co-formulated mixture, the mixture partner should provide effective disease control when used alone on the target disease and must have a different mode of action.

#### Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2024. All previous versions of this strategy are now invalid.

7. Pomefruit spray guidelines for SDHI fungicides:

Total number of spray applications per crop	1	2	3	4	5	6	7	8	9	10	11	12	>12
Max. recommended alone SDHI fungicide sprays	1	1	1	1	2	2	2	2	2	3	3	3	3
Max. recommended SDHI fungicide sprays in mixture or coformulated	1	1	2	2	2	3	3	3	3	3	3	4	4

8. **Do not** apply more than three consecutive sprays of **Group U12**, and no more than a total of six **Group U12** sprays per season.
9. If more sprays are required, tank mix **Group U12** with a protectant product at the registered rate.

**Please note:**

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2024. All previous versions of this strategy are now invalid.