

Crops(s) **Grape**

Disease(s) **Downy mildew**

Resistance management strategy for:

Group 4	Phenylamides (PA);
Group 11	Quinone outside inhibitors (QoI);
Group 11 + 3	QoI + Demethylation inhibitors (DMI);
Group 21	Quinone inside inhibitors (QiI);
Group 40	Carboxylic acid amides (CAA);
Group 45	Quinone outside inhibitor, stigmatellin binding type (QoSI);
Group 45+40	QoSI + CAA; and
Group 49	Oxysterol binding protein homologue inhibitors (OSBPI).

Guidelines:

1. Start preventative disease control sprays using **non-Group 4** protectant fungicides, typically when shoots are 10-20cm long. Continue spraying at intervals of 7-21 days depending on disease pressure, label directions and rate of vine growth.
2. **Group 4** fungicides should be applied as soon as possible after an infection period, and before the first sign of oilspots. Limit the use of **Group 4** fungicides to periods when conditions favour disease development. Always apply **Group 4** fungicides in mixtures.
3. **Group 49** fungicides should be applied prior to infection and only in mixtures with effective fungicides applied at an effective rate from a different cross resistance group. The mixing partner should give effective control of downy mildew at the rate and interval selected. A maximum of two **Group 49** applications may be made per season. Only apply **Group 49** for a maximum of one in every three sprays of the total number of downy mildew sprays. A **Group 49** application must be followed by at least two applications of a different group(s) before being reapplied.
4. Fungicide mixtures are defined as co-formulations, or tank mixes at label rate of an alternative mode of action.
5. Apply a maximum of two consecutive applications of **Group 4, 21, or 40** containing fungicides.
6. **Do not** apply **Group 11** (including **Group 11+3**) consecutively when applying alone.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2024. All previous versions of this strategy are now invalid.

7. Apply a maximum of two sprays per season of **Group 11** (including mixtures) **Group 45+40** and **Group 49**.
8. **Do not** apply a spray containing **Group 40** as the last spray of the season. Only apply a spray containing **Group 40** a maximum of 50% of the total number of downy mildew sprays.
9. Apply a maximum of three **Group 21** containing sprays per season, and a maximum of two consecutive sprays

	Group				
	4	11, 11+ 3	21+M1	40, 45+40 & 40+M3	49
Maximum number of consecutive applications	2	none	2	2	none
Maximum number of solo sprays	None	2	3	2 (50%)	None
Maximum number of sprays per season	4-mix	2	3	4-mix (50%)*	2-mix
Areas of higher agronomic risk	mix	mix	N/A	mix	mix

* Refer to points 7 &8

N.B. Consecutive sprays include mixture formulations

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2024. All previous versions of this strategy are now invalid.