

Crop(s)	Banana
Disease(s)	Yellow Sigatoka

Resistance management strategy for:

Group 3	Demethylation Inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 3	SDHI + DMI;
Group 9	Anilinopyrimidine (AP); and
Group 11	Quinone outside Inhibitor (QoI).

Guidelines

Far North Queensland

1. De-leafing must be conducted in accordance with industry guidelines.
2. Apply a regular schedule of protectant sprays.
3. When disease potential is high, apply a maximum of two consecutive **Group 3** sprays before changing to a fungicide of a different activity group.
4. **Do not** apply more than six **Group 3** sprays in any 12-month period. **Do not** apply any **Group 3** sprays in the months of June to September.
5. **Do not** apply more than two **Group 11** sprays in any 12-month period. **Do not** apply **Group 11** sprays in the months of May to September.
6. **Do not** apply more than four **Group 7** sprays in any 12-month period. **Do not** apply **Group 7** sprays in the months of June to September.
7. **Group 7** or **11** fungicides should be applied in mixture with another fungicide from a different activity Group registered for the control of Yellow Sigatoka at the full registered rate.
8. **Do not** apply consecutive sprays of **Group 7** or **11** fungicides.
9. Apply a minimum of two sprays from a different activity group between sprays of a **Group 7** or **11** fungicide.
10. **Do not** apply more than six **Group 9** sprays in any 12-month period.
11. **Do not** apply more than two consecutive sprays of a **Group 9** fungicide before changing to a fungicide of a different activity group. When using consecutive applications of **Group 9** fungicides, follow with at least as many different activity group fungicides before resuming with a **Group 9** fungicide.

Cont. on next page

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.

Group	Max. number of applications per year	Max. number of consecutive sprays	Restricted (no spray) periods
3	6 (& no more than 2 of 3 sprays)	2	June to September inclusive
7	4 (& no more than 1 of 3 sprays)	Not allowed	June to September inclusive
9	6 (& no more than 2 of 4 sprays)	2	No restriction
11	2 (& no more than 1 of 3 sprays)	Not allowed	May to September inclusive

Everywhere except far North Queensland

1. When using **Group 3** fungicides, apply a maximum of two consecutive **Group 3** sprays before changing to a fungicide of a different activity group.
2. **Do not** apply more than six **Group 3** sprays in any 12-month period.
3. **Do not** apply more than four **Group 7** or **11** sprays in any 12-month period.
4. **Do not** apply consecutive sprays of **Group 7** or **11** fungicides.
5. Apply a minimum of two sprays from a different activity group between sprays of a **Group 7** or **11** fungicide.
6. **Do not** apply more than six **Group 9** sprays in any 12-month period.
7. **Do not** apply more than two consecutive sprays of **Group 9** fungicides before changing to a fungicide of a different activity group. When using consecutive applications of **Group 9** fungicides, follow with at least as many different activity group fungicides before resuming with a **Group 9** fungicide.

Group	Max. number of applications per year	Max. number of consecutive sprays	Restricted (no spray) periods
3	6 (& no more than 2 of 3 sprays)	2	6 (& no more than 2 of 3 sprays)
7	4 (& no more than 1 of 3 sprays)	Not allowed	4 (& no more than 1 of 3 sprays)
9	6 (& no more than 2 of 4 sprays)	2	6 (& no more than 2 of 4 sprays)
11	4 (& no more than 1 of 3 sprays)	Not allowed	4 (& no more than 1 of 3 sprays)

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.