

Crop(s)	Canola
Disease(s)	Black leg and sclerotinia

Resistance management strategy for:

Group 2	Dicarboximides;
Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 3	SDHI + DMI;
Group 7 + 12	SDHI + phenylpyrroles (PP); and
Group 11 + 3	Quinone outside inhibitors + DMI

Guidelines

1. Not all listed products are registered for both diseases. Ensure the product is registered for the disease being targeted.
2. Fungicides should be used primarily as a preventative or at first sign of disease. If disease is established within the canopy, fungicides may not produce optimal results and there is very strong potential to select for fungicide resistance. Sclerotinia targeted applications should be applied during flowering of the crop, prior to an infection period. Application of fungicides for Sclerotinia may put selection pressure on the blackleg population.
3. In high risk disease environments, integrated management approaches should be used to reduce fungicide resistance risk. This includes growing canola at least 500 m from previous season's canola stubble, the use of resistant varieties, using alternative fungicide modes of action and stubble management such as knocking down and/or strategic burning.
4. The risk of developing resistance to fungicides can be reduced by incorporating different modes of action into blackleg management programs as either mixtures, co-formulations or rotations.
5. If a **Group 7** seed treatment has been used with foliar activity on blackleg (as determined by label claims), the seedling fungicide application at 4-6 leaf stage targeting blackleg should not contain a **Group 7** fungicide.
6. **Do not** apply more than two applications containing **Group 7** fungicides per growing season. Combinations of in furrow and seed treatment are counted as one application.
7. **Do not** apply more than two consecutive applications of a **Group 3** fungicide.
8. **Do not** apply more than one application containing a **Group 11** fungicide.
9. Minimise use of fungicides which are known to have compromised resistance status.
10. If seasonal conditions require a second fungicide application at 50% flowering after a 20% flowering timing, the second application should be from a different **Group**.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.

Application stage (Disease being controlled)		Rotation options for difference fungicide active groups											
Seed dressing & in-furrow (Blackleg)		None	None	None	None	None	None	3	3	3	3	3	3
Seedling foliar (Blackleg)		None	3	7	7+3	7+12	11+3	None	3	7	7+3	7+12	11+3
20% flower (Sclerotinia) Choose only one option from this section	1	None	None	None	None	None	None	None	None	None	None	None	None
	2	2	2	2	2	2	2	2	2	2	2	2	2
	3	3	3	3	3	3	3		3		3	3	
	4	7+3	7+3	7+3	7+3	7+3			7+3		7+3	7+3	
	5	7+12	7+12	7+12	7+12		7+12	7+12	7+12	7+12			7+12
	6	11+3	11+3	11+3	11+3	11+3			11+3		11+3	11+3	

Application stage (Disease being controlled)		Rotation options for difference fungicide active groups					
Seed dressing & in-furrow (Blackleg)		7	7	7	7 + 3	7 + 3	7 + 3
Seedling foliar (Blackleg)		None	3	11+3	None	3	11+3
20% flower (Sclerotinia) Choose only one option from this section	1	None	None	None	None	None	None
	2	2	2	2	2	2	2
	3	3	3	3	3		
	4	7+3	7+3	7+3	7+3		
	5	7+12	7+12	7+12	7+12	7+12	7+12
	6	11+3	11+3		11+3		

If a second application at 50% flower required:

		Application at 20% flowering				
		2	3	7+3	7+12	11+3
Rotation options for 50% flowering second application			2	2	2	2
	3				3	
	7+3					
	11+3				11+3	

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.