

Crop(s) **Macadamia**
Disease(s) **Husk spot**

Resistance management strategy for:

Group 1	Methyl benzimidazole carbamates (MBC);
Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitor (SDHI);
Group 7 + 3	SDHI + DMI
Group 11	Quinone outside inhibitors (QoI);
Group 7 + 11	SDHI + QoI; and
Group 11 + 3	QoI + DMI.

Guidelines

		Max. recommended number of sprays which can contain Group:						
		1	3	7	11	7+3	7+11	11+3
Total number of husk spot targeting sprays	1	1	1	1	1	1	1	1
	2	2	2	1	1	2	2	2
	3	2	2	1	1	2	2	2
	4+	2	2	1	1	2	2	2

1. If application of a single group product is used, **do not** apply consecutive sprays of the same group.
2. If applying a pre-mixture or tank mixture product containing two different groups, **do not** apply more than two consecutive sprays before changing to another Mode of Action group. This includes from one season to the next, and applications made to manage other diseases.
3. The spray program should be considered, and the strategy applied on a whole-orchard basis.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.