

Crop(s) Potato

Insect(s) Potato tuber moth / Tomato leafminer (Phthorimaea

opercululla)

## Guidelines

1. Monitor pest levels and **do not** spray unless pest thresholds are exceeded.\

- 2. **Rotate** insecticide groups and **do not** apply more than two consecutive applications of products with the same Mode of Action.
- 3. Integrate both chemical and non-chemical means of control as part of the overall control strategy. Examples are the use of predators/parasites and relevant cultural practices (crop hygiene, rotation of planted areas, and strategic time of planting).

Group*	Chemical subgroup	Example chemical
1A	Carbamates	carbaryl, methomyl
1B	Organophosphates	acephate, azinphos-methyl, methamidophos, diazinon, dichlorvos
3A	Pyrethroids	permethrin
5	Spinosyns	spinosad, spinetoram
28	Diamides	chlorantraniliprole, flubendiamide, cyantraniliprole

<sup>\*</sup> Refer: CropLife Australia Expert Committee on Insecticide Resistance Mode of Action Classification for Insecticides.

## Notes regarding the application of insecticides:

To ensure the most effective control of the pest:

- a) Always read and follow product labels and use the full recommended label rates of application; and
- b) Ensure the spray equipment is properly calibrated and achieves good coverage of target area with appropriately sized spray droplets.